

Wiener index of graphs in terms of eccentricities

H. S. RAMANE¹, A. B. GANAGI^{2,*}, H. B. WALIKAR³

¹Department of Mathematics, Karnatak University, Dharwad – 580003, India

²Department of Mathematics, Gogte Institute of Technology, Udyambag, Belgaum–590008, India

³Department of Computer Science, Karnatak University, Dharwad – 580003, India

(Received August 15, 2013; Accepted October 26, 2013)

ABSTRACT

The Wiener index $W(G)$ of a connected graph G is defined as the sum of the distances between all unordered pairs of vertices of G . The eccentricity of a vertex v in G is the distance to a vertex farthest from v . In this paper we obtain the Wiener index of a graph in terms of eccentricities. Further we extend these results to the self-centered graphs.

Keywords: Wiener index, distance, eccentricity, radius, diameter, self-centered graph.

1. INTRODUCTION

The Wiener index $W(G)$ of a connected graph G is defined as the sum the distances between all unordered pairs of vertices of G . It was put forward by Harold Wiener [1]. The Wiener index is a graph invariant intensively studied both in mathematics and chemical literature. For details one may refer [2 – 13] and the reference cited there in.

Let G be a connected, simple graph with vertex set $V(G)$. The degree of a vertex v in G is the number of edges incident to it and is denoted by $deg(v)$. The distance between the vertices u and v , denoted by $d(u,v)$, is the length of the shortest path joining them. The eccentricity $e(v)$ of a vertex v is the distance to a vertex farthest from v , that is

$$e(v) = \max\{d(u,v) \mid u \in V(G)\}.$$

The radius $r(G)$ of a graph G is the minimum eccentricity of the vertices and the diameter $d(G)$ of G is the maximum eccentricity. A vertex v is called central vertex of G if $e(v) = r(G)$. A graph is called self-centered if every vertex is a central vertex. Thus in a self-centered graph $r(G) = d(G)$. An eccentric vertex of a vertex v is a vertex farthest away from v . An eccentric path of a vertex v denoted by $P(v)$ is a path of length $e(v)$ joining v and its eccentric vertex. There may exists more than one eccentric path for a given vertex.

*Corresponding author (Email: abganagi@yahoo.co.in)

If v_1, v_2, \dots, v_n are the vertices of graph G then the Wiener index of G is defined as

$$W(G) = \sum_{1 \leq i < j \leq n} d(v_i, v_j).$$

The distance number of a vertex v_i of a graph G denoted by $d(v_i | G)$ is defined as

$$d(v_i | G) = \sum_{j=1}^n d(v_i, v_j).$$

Therefore

$$W(G) = \frac{1}{2} \sum_{i=1}^n d(v_i | G).$$

In this paper we obtain the Wiener index in terms of eccentricities. For graph theoretic terminology we refer the book [14].

2. MAIN RESULTS

Theorem 2.1: Let G be a connected graph with n vertices, m edges and $e_i = e(v_i)$, $i = 1, 2, \dots, n$, then

$$W(G) \geq \frac{1}{2} \left[n(2n-1) - 2m + \sum_{i=1}^n \frac{e_i(e_i-3)}{2} \right]. \quad (1)$$

Equality holds if and only if for every vertex v_i of G , if $P(v_i)$ is one of the eccentric path of v_i , then for every $v_j \in V(G)$ which is not on $P(v_i)$, $d(v_i, v_j) \leq 2$.

Proof: Let $P(v_i)$ be one of the eccentric path of $v_i \in V(G)$.

Let $A_1(v_i) = \{v_j | v_j \text{ is on eccentric path } P(v_i) \text{ of } v_i\}$,

$A_2(v_i) = \{v_j | v_j \text{ is adjacent to } v_i \text{ and which is not on the eccentric path } P(v_i) \text{ of } v_i\}$,

$A_3(v_i) = \{v_j | v_j \text{ is not adjacent to } v_i \text{ and not on the eccentric path } P(v_i) \text{ of } v_i\}$.

Clearly $A_1(v_i) \cup A_2(v_i) \cup A_3(v_i) = V(G)$ and

$$|A_1(v_i)| = e_i + 1, \quad |A_2(v_i)| = \deg(v_i) - 1, \quad |A_3(v_i)| = n - e_i - \deg(v_i).$$

$$\text{Now } \sum_{v_j \in A_1(v_i)} d(v_i, v_j) = 1 + 2 + \dots + e_i = \frac{e_i(e_i+1)}{2},$$

$$\sum_{v_j \in A_2(v_i)} d(v_i, v_j) = \deg(v_i) - 1,$$

$$\sum_{v_j \in A_3(v_i)} d(v_i, v_j) \geq 2(n - e_i - \deg(v_i)).$$

Therefore,

$$\begin{aligned}
 d(v_i | G) &= \sum_{j=1}^n d(v_i, v_j) \\
 &= \sum_{v_j \in A_1(v_i)} d(v_i, v_j) + \sum_{v_j \in A_2(v_i)} d(v_i, v_j) + \sum_{v_j \in A_3(v_i)} d(v_i, v_j) \\
 &\geq \frac{e_i(e_i + 1)}{2} + \deg(v_i) - 1 + 2(n - e_i - \deg(v_i)) \\
 &= 2n - \deg(v_i) - 1 + \frac{e_i(e_i - 3)}{2}.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i | G) \\
 &\geq \frac{1}{2} \sum_{i=1}^n \left[2n - \deg(v_i) - 1 + \frac{e_i(e_i - 3)}{2} \right] \\
 &= \frac{1}{2} \left[2n^2 - 2m - n + \sum_{i=1}^n \frac{e_i(e_i - 3)}{2} \right] \\
 &= \frac{1}{2} \left[n(2n - 1) - 2m + \sum_{i=1}^n \frac{e_i(e_i - 3)}{2} \right].
 \end{aligned}$$

For equality,

Let G be a graph and $P(v_i)$ be one of the eccentric paths of $v_i \in V(G)$. Let $A_1(v_i)$, $A_2(v_i)$ and $A_3(v_i)$ be the sets as defined in the first part of the proof of this theorem.

Let $d(v_i, v_j) = 2$, where $v_j \in A_3(v_i)$.

Therefore $\sum_{v_j \in A_3(v_i)} d(v_i, v_j) = 2(n - e_i - \deg(v_i))$,

$$\sum_{v_j \in A_1(v_i)} d(v_i, v_j) = \frac{e_i(e_i + 1)}{2} \quad \text{and} \quad \sum_{v_j \in A_2(v_i)} d(v_i, v_j) = \deg(v_i) - 1$$

Thus

$$\begin{aligned}
 d(v_i | G) &= \sum_{j=1}^n d(v_i, v_j) \\
 &= \sum_{v_j \in A_1(v_i)} d(v_i, v_j) + \sum_{v_j \in A_2(v_i)} d(v_i, v_j) + \sum_{v_j \in A_3(v_i)} d(v_i, v_j) \\
 &= \frac{e_i(e_i + 1)}{2} + \deg(v_i) - 1 + 2(n - e_i - \deg(v_i)) \\
 &= 2n - \deg(v_i) - 1 + \frac{e_i(e_i - 3)}{2}.
 \end{aligned}$$

Hence

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i | G) \\
&= \frac{1}{2} \sum_{i=1}^n \left[2n - \deg(v_i) - 1 + \frac{e_i(e_i - 3)}{2} \right] \\
&= \frac{1}{2} \left[2n^2 - 2m - n + \sum_{i=1}^n \frac{e_i(e_i - 3)}{2} \right] \\
&= \frac{1}{2} \left[n(2n - 1) - 2m + \sum_{i=1}^n \frac{e_i(e_i - 3)}{2} \right].
\end{aligned}$$

Conversely,

Suppose G is not such graph as defined in the equality part of this theorem. Then there exist at least one vertex $v_j \in A_3(v_i)$ such that $d(v_i, v_j) \geq 3$. Let $A_3(v_i)$ be partitioned into two sets $A_{31}(v_i)$ and $A_{32}(v_i)$, where

$$A_{31}(v_i) = \{v_j \mid v_j \text{ is not adjacent to } v_i, \text{ not on the eccentric path } P(v_i) \text{ of } v_i \text{ and } d(v_i, v_j) = 2\}$$

$$A_{32}(v_i) = \{v_j \mid v_j \text{ is not adjacent to } v_i, \text{ not on the eccentric path } P(v_i) \text{ of } v_i \text{ and } d(v_i, v_j) \geq 3\}.$$

Let $|A_{32}(v_i)| = l \geq 1$. So, $|A_{31}(v_i)| = n - e_i - \deg(v_i) - l$.

$$\text{Therefore } \sum_{v_j \in A_1(v_i)} d(v_i, v_j) = \frac{e_i(e_i + 1)}{2}, \quad \sum_{v_j \in A_2(v_i)} d(v_i, v_j) = \deg(v_i) - 1,$$

$$\sum_{v_j \in A_{31}(v_i)} d(v_i, v_j) = 2(n - e_i - \deg(v_i) - l) \text{ and } \sum_{v_j \in A_{32}(v_i)} d(v_i, v_j) \geq 3l.$$

Therefore

$$\begin{aligned}
d(v_i | G) &= \sum_{j=1}^n d(v_i, v_j) \\
&= \sum_{v_j \in A_1(v_i)} d(v_i, v_j) + \sum_{v_j \in A_2(v_i)} d(v_i, v_j) + \sum_{v_j \in A_{31}(v_i)} d(v_i, v_j) + \sum_{v_j \in A_{32}(v_i)} d(v_i, v_j) \\
&\geq \frac{e_i(e_i + 1)}{2} + \deg(v_i) - 1 + 2(n - e_i - \deg(v_i) - l) + 3l \\
&= 2n - \deg(v_i) - 1 + \frac{e_i(e_i - 3)}{2} + l.
\end{aligned}$$

Therefore

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i | G) \\
&\geq \frac{1}{2} \sum_{i=1}^n \left[2n - \deg(v_i) - 1 + \frac{e_i(e_i - 3)}{2} + l \right] \\
&= \frac{1}{2} \left[2n^2 - 2m - n + \sum_{i=1}^n \frac{e_i(e_i - 3)}{2} + nl \right]
\end{aligned}$$

$$\geq \frac{1}{2} \left[n(2n-1) - 2m + \sum_{i=1}^n \frac{e_i(e_i-3)}{2} \right] \text{ as } l \geq 1, \text{ which is a contradiction.}$$

This contradiction proves the result. \square

Corollary 2.2: Let G be a self-centered graph with n vertices, m edges and radius $r = r(G)$, then $W(G) \geq \frac{1}{2} \left[n(2n-1) - 2m + \frac{nr(r-3)}{2} \right]$.

Equality holds if and only if for every vertex v_i of a self-centered graph G , if $P(v_i)$ is one of the eccentric path of v_i then for every $v_j \in V(G)$ which is not on the eccentric path $P(v_i)$, $d(v_i, v_j) \leq 2$.

Proof: For self-centered graph each vertex has same eccentricity equal to the radius r , that is, $e_i = e(v_i) = r, i = 1, 2, \dots, n$. Therefore from Eq. (1)

$$\begin{aligned} W(G) &\geq \frac{1}{2} \left[n(2n-1) - 2m + \sum_{i=1}^n \frac{r(r-3)}{2} \right] \\ &= \frac{1}{2} \left[n(2n-1) - 2m + \frac{nr(r-3)}{2} \right] \end{aligned}$$

The proof of the equality part is similar to the proof of equality part of Theorem 1.1. \square

Theorem 2.3: Let G be a connected graph with n vertices and $e_i = e(v_i), i = 1, 2, \dots, n$, then

$$W(G) \geq \frac{1}{2} \left[n^2 + \sum_{i=1}^n \frac{(e_i+1)(e_i-2)}{2} \right]. \tag{2}$$

Equality holds if and only if for every vertex v_i of G , if $P(v_i)$ is one of the eccentric path of v_i , then for every $v_j \in V(G)$ which is not on $P(v_i)$, $d(v_i, v_j) = 1$.

Proof: Let $e_i = e(v_i), i = 1, 2, \dots, n$ and $P(v_i)$ be one of the eccentric path of $v_i \in V(G)$.

Let $B_1(v_i) = \{v_j \mid v_j \text{ is on eccentric path } P(v_i) \text{ of } v_i\}$,

$B_2(v_i) = \{v_j \mid v_j \text{ is not on the eccentric path } P(v_i) \text{ of } v_i\}$.

Clearly $B_1(v_i) \cup B_2(v_i) = V(G)$ and

$$|B_1(v_i)| = e_i + 1, \quad |B_2(v_i)| = n - e_i - 1.$$

Now $\sum_{v_j \in B_1(v_i)} d(v_i, v_j) = 1 + 2 + \dots + e_i = \frac{e_i(e_i+1)}{2}$,

$$\sum_{v_j \in B_2(v_i)} d(v_i, v_j) \geq 1(n - e_i - 1),$$

Therefore

$$\begin{aligned}
d(v_i | G) &= \sum_{j=1}^n d(v_i, v_j) \\
&= \sum_{v_j \in B_1(v_i)} d(v_i, v_j) + \sum_{v_j \in B_2(v_i)} d(v_i, v_j) \\
&\geq \frac{e_i(e_i + 1)}{2} + n - e_i - 1 \\
&= n + \frac{(e_i - 2)(e_i + 1)}{2}.
\end{aligned}$$

Therefore

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i | G) \\
&\geq \frac{1}{2} \sum_{i=1}^n \left[n + \frac{(e_i - 2)(e_i + 1)}{2} \right] \\
&= \frac{1}{2} \left[n^2 + \sum_{i=1}^n \frac{(e_i - 2)(e_i + 1)}{2} \right].
\end{aligned}$$

For equality,

Let G be a graph and $P(v_i)$ be one of the eccentric paths of $v_i \in V(G)$. Let $B_1(v_i)$ and $B_2(v_i)$ be the sets as defined in the first part of the proof of this theorem.

Let $d(v_i, v_j) = 1$, where $v_j \in B_2(v_i)$.

Therefore $\sum_{v_j \in B_2(v_i)} d(v_i, v_j) = n - e_i - 1$ and $\sum_{v_j \in B_1(v_i)} d(v_i, v_j) = \frac{e_i(e_i + 1)}{2}$.

Therefore

$$\begin{aligned}
d(v_i | G) &= \sum_{j=1}^n d(v_i, v_j) \\
&= \sum_{v_j \in B_1(v_i)} d(v_i, v_j) + \sum_{v_j \in B_2(v_i)} d(v_i, v_j) \\
&= \frac{e_i(e_i + 1)}{2} + n - e_i - 1 \\
&= n + \frac{(e_i - 2)(e_i + 1)}{2}.
\end{aligned}$$

Therefore

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i | G) \\
&= \frac{1}{2} \sum_{i=1}^n \left[n + \frac{(e_i - 2)(e_i + 1)}{2} \right]
\end{aligned}$$

$$= \frac{1}{2} \left[n^2 + \sum_{i=1}^n \frac{(e_i - 2)(e_i + 1)}{2} \right].$$

Conversely,

Suppose G is not such graph as defined in the equality part of this theorem. Then there exist at least one vertex $v_j \in B_2(v_i)$ such that $d(v_i, v_j) \geq 2$. Let $B_2(v_i)$ be partitioned into two sets $B_{21}(v_i)$ and $B_{22}(v_i)$, where

$$B_{21}(v_i) = \{v_j \mid v_j \text{ is not on the eccentric path } P(v_i) \text{ of } v_i \text{ and } d(v_i, v_j) = 1\}$$

$$B_{22}(v_i) = \{v_j \mid v_j \text{ is not on the eccentric path } P(v_i) \text{ of } v_i \text{ and } d(v_i, v_j) \geq 2\}.$$

Let $|B_{22}(v_i)| = l \geq 1$

Therefore $|B_{21}(v_i)| = n - e_i - 1 - l$.

Therefore $\sum_{v_j \in B_1(v_i)} d(v_i, v_j) = \frac{e_i(e_i + 1)}{2}$, $\sum_{v_j \in B_{21}(v_i)} d(v_i, v_j) = n - e_i - 1 - l$ and $\sum_{v_j \in B_{22}(v_i)} d(v_i, v_j) \geq 2l$.

Therefore

$$\begin{aligned} d(v_i \mid G) &= \sum_{j=1}^n d(v_i, v_j) \\ &= \sum_{v_j \in B_1(v_i)} d(v_i, v_j) + \sum_{v_j \in B_{21}(v_i)} d(v_i, v_j) + \sum_{v_j \in B_{22}(v_i)} d(v_i, v_j) \\ &\geq \frac{e_i(e_i + 1)}{2} + n - e_i - 1 - l + 2l \\ &= n + l + \frac{(e_i - 2)(e_i + 1)}{2}. \end{aligned}$$

Therefore

$$\begin{aligned} W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i \mid G) \\ &\geq \frac{1}{2} \sum_{i=1}^n \left[n + l + \frac{(e_i - 2)(e_i + 1)}{2} \right] \\ &\geq \frac{1}{2} \sum_{i=1}^n \left[n + 1 + \frac{(e_i - 2)(e_i + 1)}{2} \right] \text{ as } l \geq 1. \\ &= \frac{1}{2} \left[n(n + 1) + \sum_{i=1}^n \frac{(e_i - 2)(e_i + 1)}{2} \right]. \end{aligned}$$

This is a contradiction. Hence the proof. □

If G is a self-centered graph then $e_i = e(v_i) = r(G)$ for all $i = 1, 2, \dots, n$. Substituting this in Eq. (2) we get following corollary.

Corollary 2.4: Let G be a self-centered graph with n vertices and radius $r = r(G)$, then

$$W(G) \geq \frac{1}{2} \left[n^2 + \frac{n(r+1)(r-2)}{2} \right].$$

Equality holds if and only if for every vertex v_i of a self-centered graph G , if $P(v_i)$ is one of the eccentric path of v_i then for every $v_j \in V(G)$ which is not on the eccentric path $P(v_i)$, $d(v_i, v_j) = 1$.

Theorem 2.5: Let G be a connected graph with n vertices, m edges and $diam(G) = d$. Let $e_i = e(v_i)$, $i = 1, 2, \dots, n$, then

$$W(G) \leq \frac{1}{2} \left[n(nd - 1) - (1 - d)2m + \sum_{i=1}^n \frac{e_i(e_i + 1 - 2d)}{2} \right]. \quad (3)$$

Equality holds if and only if $diam(G) \leq 2$.

Proof: Let $P(v_i)$ be one of the eccentric path of $v_i \in V(G)$.

Let $A_1(v_i) = \{v_j \mid v_j \text{ is on the eccentric path } P(v_i) \text{ of } v_i\}$,

$A_2(v_i) = \{v_j \mid v_j \text{ is adjacent to } v_i \text{ and which is not on the eccentric path } P(v_i) \text{ of } v_i\}$,

$A_3(v_i) = \{v_j \mid v_j \text{ is not adjacent to } v_i \text{ and not on the eccentric path } P(v_i) \text{ of } v_i\}$.

Clearly $A_1(v_i) \cup A_2(v_i) \cup A_3(v_i) = V(G)$ and

$$|A_1(v_i)| = e_i + 1, \quad |A_2(v_i)| = deg(v_i) - 1, \quad |A_3(v_i)| = n - e_i - deg(v_i).$$

Now
$$\sum_{v_j \in A_1(v_i)} d(v_i, v_j) = 1 + 2 + \dots + e_i = \frac{e_i(e_i + 1)}{2},$$

$$\sum_{v_j \in A_2(v_i)} d(v_i, v_j) = deg(v_i) - 1,$$

$$\sum_{v_j \in A_3(v_i)} d(v_i, v_j) \leq d(n - e_i - deg(v_i)).$$

Therefore

$$\begin{aligned} d(v_i \mid G) &= \sum_{j=1}^n d(v_i, v_j) \\ &= \sum_{v_j \in A_1(v_i)} d(v_i, v_j) + \sum_{v_j \in A_2(v_i)} d(v_i, v_j) + \sum_{v_j \in A_3(v_i)} d(v_i, v_j) \\ &\leq \frac{e_i(e_i + 1)}{2} + deg(v_i) - 1 + d(n - e_i - deg(v_i)) \\ &= nd - 1 + (1 - d)deg(v_i) + \frac{e_i(e_i + 1 - 2d)}{2}. \end{aligned}$$

Therefore

$$\begin{aligned}
 W(G) &= \frac{1}{2} \sum_{i=1}^n d(v_i | G) \\
 &\leq \frac{1}{2} \sum_{i=1}^n \left[nd - 1 + (1-d)deg(v_i) + \frac{e_i(e_i + 1 - 2d)}{2} \right] \\
 &= \frac{1}{2} \left[n(nd - 1) + (1-d)2m + \sum_{i=1}^n \frac{e_i(e_i + 1 - 2d)}{2} \right] \quad \text{since} \\
 &\quad \sum_{i=1}^n deg(v_i) = 2m.
 \end{aligned}$$

For equality,

Let $diam(G) \leq 2$.

Case 1: If $diam(G) = 1$ then $G = K_n$. Therefore $A_3(v_i) = \Phi$ and $e_i = e(v_i) = 1, i = 1, 2, \dots, n$.

Therefore $W(G) = \frac{1}{2} \left[n(n-1) + \sum_{i=1}^n \frac{1(1+1-2)}{2} \right] = \frac{n(n-1)}{2}$.

Case 2: If $diam(G) = 2$, then for $v_j \in A_3(v_i), d(v_i, v_j) = 2$.

Therefore $\sum_{v_j \in A_3(v_i)} d(v_i, v_j) = 2(n - e_i - deg(v_i))$.

Hence $W(G) = \frac{1}{2} \left[n(nd - 1) + (1-d)2m + \sum_{i=1}^n \frac{e_i(e_i + 1 - 2d)}{2} \right]$
 $= \frac{1}{2} \left[n(2n - 1) - 2m + \sum_{i=1}^n \frac{e_i(e_i - 3)}{2} \right]$.

Conversely,

$$\begin{aligned}
 d(v_i | G) &= \sum_{j=1}^n d(v_i, v_j) \\
 &= \sum_{v_j \in A_1(v_i)} d(v_i, v_j) + \sum_{v_j \in A_2(v_i)} d(v_i, v_j) + \sum_{v_j \in A_3(v_i)} d(v_i, v_j)
 \end{aligned} \tag{4}$$

The first summation of Eq. (4) contains the distance between v_i and the vertices on its eccentric path $P(v_i)$. Second summation of Eq. (4) contains the distance between v_i and its neighbor which are not on the eccentric path $P(v_i)$. The third summation of Eq. (4) contains the distance between v_i and a vertex which is neither adjacent to v_i nor on the eccentric path $P(v_i)$. Hence the equality in Eq. (4) holds if and only if $d = diam(G) \leq 2$. It is true for all $v_i \in V(G)$. Hence $diam(G) \leq 2$. □

Corollary 2.6: Let G be a self-centered graph with n vertices and radius $r = r(G)$, then

$$W(G) \leq \frac{1}{2} \left[n(nr - 1) - \frac{(r-1)(nr + 4m)}{2} \right].$$

Equality holds if and only if $diam(G) \leq 2$.

Proof: Proof follows by substituting $e_i = e(v_i) = r$, $i = 1, 2, \dots, n$ in Eq. (3). \square

REFERENCES

1. H. Wiener, Structural determination of paraffin boiling points, *J. Amer. Chem. Soc.*, **69** (1947), 17 – 20.
2. F. Buckley, F. Harary, *Distances in Graphs*, Addison-Wesley, Redwood, 1990.
3. I. Gutman, Y. N. Yeh, S. L. Lee, Y. L. Luo, Some recent results in the Theory of the Wiener number, *Indian J. Chem.*, **32A** (1993), 651 – 661.
4. R. C. Entringer, Distance in graphs: Trees, *J. Combin. Math. Combin. Comput.*, **24** (1997), 65 – 84.
5. A. A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: Theory and Applications, *Acta Appl. Math.*, **66** (2001), 211 – 249.
6. A. Dobrynin, I. Gutman, S. Klavzar, P. Zigert, Wiener index of hexagonal systems, *Acta Appl. Math.*, **72**(2002), 247 – 294.
7. I. Gutman, G. Zenkevich, Wiener index and vibrational energy, *Z. Naturforsch.*, **57A**(2002), 824 – 828.
8. H. B. Walikar, H. S. Ramane, V. S. Shigehalli, Wiener number of Dendrimers, In: *Proc. National Conf. on Mathematical and Computational Models*, (Eds. R. Nadarajan and G. Arulmozhi), Allied Publishers, New Delhi, 2003, pp. 361 – 368.
9. H. B. Walikar, V. S. Shigehalli, H. S. Ramane, Bounds on the Wiener number of a graph, *MATCH Commun. Math. Comp. Chem.*, **50** (2004), 117 – 132.
10. G. C. Garcia, I. L. Ruiz, M. A. Gomez-Nieto, J. A. Doncel, A. G. Plaza, From Wiener index to molecule, *J. Chem. Inf. Model.*, **45** (2005), 231 – 238.
11. H. Liu, X. F. Pan, On the Wiener index of trees with fixed diameter, *MATCH Commun. Math. Comput. Chem.*, **60** (2008), 85 – 94.
12. S. Wang, X. Guo, Trees with extremal Wiener indices, *MATCH Commun. Math. Comput. Chem.*, **60** (2008), 609 – 622.
13. A. Chon, F. Zhang, Wiener index and perfect matching in random phenylene chains, *MATCH Commun. Math. Comput. Chem.*, **61** (2009), 623 – 630.
14. K. C. Das, I. Gutman, Estimating the Wiener index by means of number of vertices of edges and diameter, *MATCH Commun. Math. Comput. Chem.*, **64** (2010), 647 – 660.